

The impact of prescribed fire versus wildfire on the immune and cardiovascular systems of children

To the Editor

The increase in wildfires associated with climate change augments the impact of air pollution on health in many areas of the country. When wildfires occur, there is an increase in asthma attacks and associated comorbidities,^{1,2} especially for asthma hospitalization in ages 0-5 years³ and more recently, it has been shown that there are increases in cardiovascular events.⁴ Given the health risks associated with high-intensity wildfires, there is motivation to increase the use of lower intensity prescribed fires. Prescribed burns decrease the buildup of flammable vegetation and subsequent fuel for wildfires, mitigating the spread and intensity of wildfires. However, prescribed fire raises public concerns because of the additional pollutant exposure.

Therefore, our objective is to determine whether there are differential health consequences with a prescribed fire vs wildfire. We focus on children given their reduced lung size, increased metabolic rates, higher respiratory rate, and developing immune systems,⁵ and because in macaque monkeys who are exposed to wildfire smoke in infancy, there is associated immune dysregulation and decreased lung function in adolescence.⁶ We hypothesize that the health impacts of a prescribed fire are less detrimental to the respiratory and cardiovascular systems than a wildfire in school-aged children and that T-cell skewing and epigenetic modulation will occur with exposure to wildfire more than from exposure to a prescribed fire.

We analyzed data collected from a convenience sample of subjects (n = 220) over a period of 2 years living in Fresno, CA, all of whom were potentially exposed to smoke from fires, which consisted of similar varieties of coniferous trees, in nearby Yosemite National Park. Health questionnaires, blood samples, and vital signs were collected, and subjects were selected that had their blood drawn 3 months after a prescribed fire or wildfire, because our prior research indicates that this time frame is associated with increased methylation of the *Foxp3* gene.⁷ Using this criteria, we analyzed data from 32 children (median age = 7 [range 7; 8] yrs, 38% asthmatic as per NHLBI guidelines) exposed to a prescribed fire 70 miles away covering 553 acres in March, 2015, and 36 children (median age = 8 [range 7; 8] yrs, 25% asthmatic) exposed to a wildfire 70 miles away covering 415 acres in September 2015. A control group of 18 children was also compared (median age = 8 [range 7; 8] yrs; 21% asthmatic), who had no obvious exposure to wildfires or prescribed fire and were living in the San Francisco Bay area, where pollution levels

are consistently low (ie, <10 $\mu\text{g}/\text{m}^3$ of $\text{PM}_{2.5}$). All subjects were consented with an IRB-approved protocol.

Pollution exposure was measured (4, 5, and 6-ringed polycyclic aromatic hydrocarbons (PAH_{456}), particulate matter with aerodynamic diameter of 2.5 μm or less ($\text{PM}_{2.5}$), particulate matter with aerodynamic diameter of 10 μm or less (PM_{10}), elemental carbon (EC), ozone (O_3), carbon monoxide (CO), nitrogen dioxide (NO_2), and nitrogen oxides (NO_x)) from four central site monitors and both distance-weighted to the subject's home as in previous studies⁸ and averaged across the monitoring sites. Peripheral blood mononuclear cells were stained with metal-conjugated antibodies for surface markers, and CyTOF was performed. Methylation studies using pyrosequencing were performed per published methods⁶ on selected CpG sites of the *Foxp3*, *IL-4*, *IL-10*, and *IFN γ* genes.

As shown in Figure 1, all pollutant exposures were higher in the wildfire group (n = 36) than the prescribed fire (n = 32) groups ($P < 0.0001$; wildfire vs prescribed means: $\text{NO}_2 = 10.7$ parts per billion [ppb] ± 0.3 vs 4.0 ppb ± 0.2 ; $\text{NO}_x = 25.6$ ppb ± 1.0 vs 9.9 ppb ± 0.5 ; $\text{PAH}_{456} = 11.4$ ng/ $\text{m}^3 \pm 0.4$ vs 5.3 ng/ $\text{m}^3 \pm 0.2$; EC = 1.0 $\mu\text{g}/\text{m}^3 \pm 0.02$ vs 0.48 $\mu\text{g}/\text{m}^3 \pm 0.01$; CO = 0.56 parts per million [ppm] ± 0.02 vs 0.25 ppm ± 0.01 ; $\text{PM}_{10} = 41.5$ $\mu\text{g}/\text{m}^3 \pm 1.1$ vs 28.0 $\mu\text{g}/\text{m}^3 \pm 0.3$; $\text{PM}_{2.5} = 15.9$ $\mu\text{g}/\text{m}^3 \pm 0.4$ vs 10.0 $\mu\text{g}/\text{m}^3 \pm 0.2$). In addition, average $\text{PM}_{2.5}$ levels were calculated 2 weeks prior to each fire, throughout each fire and 2 weeks after each fire to determine the potential contributions of each fire. $\text{PM}_{2.5}$ levels increased during the wildfire and then returned to baseline indicating that the wildfire was likely associated with the rise in $\text{PM}_{2.5}$ levels (2 weeks prior mean = 9.3 $\mu\text{g}/\text{m}^3$ [SD = 2.5]; during fire mean = 13.7 $\mu\text{g}/\text{m}^3$ [SD = 5.7] vs 2 weeks after mean = 9.1 $\mu\text{g}/\text{m}^3$ [SD = 1.9]). For the prescribed fire, the $\text{PM}_{2.5}$ levels decreased 12 $\mu\text{g}/\text{m}^3$ from prefire to postfire, indicating that the prescribed burn likely did not contribute substantially to $\text{PM}_{2.5}$ levels (2 weeks prior mean = 17.8 $\mu\text{g}/\text{m}^3$ [SD = 5.9]; duration of fire mean = 8.5 $\mu\text{g}/\text{m}^3$ [SD = 3.5]; 2 weeks post mean = 5.8 $\mu\text{g}/\text{m}^3$ [SD = 2.4]).

To investigate the immune system, immunophenotype results were compared with a one-way ANOVA across the 3 groups for percent Th1 cells (CD4+, CXCR3+, CCR5+), revealing significant differences among groups ($P < 0.0001$) as shown in Figure 2, with the lowest Th1% for the wildfire group (control $5.19\% \pm 1.89$; prescribed fire $3.99\% \pm 0.34$; wildfire $2.04\% \pm 0.31$). There were no significant differences between the groups for other immune cell types, such as Th2 cells (CD4+, CCR4+, CCR6-; $P = 0.14$) or T regulatory cells (CD4+, CD 25+, CD 127-; $P = 0.66$).




FIGURE 1 Average levels of pollutants during the wildfire and prescribed fire. When comparing prescribed vs wildfire, $P < 0.0001$ for each pollutant shown




FIGURE 2 Th1 Cell percentage of CD 4+ cells for children 90 d after being exposed to a prescribed fire, wildfire, or no exposure (1-way ANOVA, $P < 0.0001$)

Methylation levels between the prescribed and wildfire groups were compared using linear regression models while controlling for covariates (age, sex, BMI percentile, race, second-hand smoke, and asthma status). Foxp3 methylation in the promoter region of DNA isolated from the same blood samples was increased post wildfire exposure compared to prescribed fire exposure (B estimate [est] = 2.59; Standard error [SE] = 0.95; $P = 0.01$). Moreover, there was a trend toward worsened health outcomes in the wildfire group compared to the prescribed group, including increases in wheezing episodes in those with no prior history of asthma, increases in asthma exacerbations in those with prior asthma, and rises in pulse pressure (est = 4.08; SE = 2.35; $P = 0.09$).

The increase in Foxp3 methylation associated with the wildfire is consistent with prior air pollution studies.^{7,8} The reduction in Th1 pro-inflammatory T cells associated with wildfire exposure may be consistent with the molecular heterogeneity of asthma and associated endotypes.⁹ Moreover, in cardiovascular disease, which is an inflammatory process and also associated with air pollution

and wildfires, Th1 cells have been associated with immunity in atherosclerosis.¹⁰ While this is a descriptive, retrospective study and the PM levels do not distinguish from various sources including fires, these preliminary results suggest future studies are needed. This will allow us to both understand the mechanism by which wildfire exposure impacts the immune system and to investigate the health impact of prescribed fire vs wildfire, as there is heightened motivation to increase the application of prescribed burns to combat the risks of increasing wildfire size and intensity in several areas of the country.

CONFLICT OF INTEREST

Dr Kari Nadeau indicates that she has received funding, is currently funded by, or has cofounded the following: National Institutes of Health (NIH), Food Allergy Research & Education (FARE), End Allergies Together (EAT), Before Brands, Alladapt Immunotherapeutics, Adare Pharmaceuticals, AstraZeneca, Novartis, Genentech, Astellas, DBV Technologies, ForTra, Aimmune Therapeutics, Regeneron, Sanofi, Nestle, and the Environmental Protection Agency (EPA).

FUNDING INFORMATION

The Nature Conservancy, NIEHS, NHLBI, and Sean N Parker Center for Allergy and Asthma Research at Stanford University.

Mary Prunicki¹ 
 Rodd Kelsey²
 Justin Lee¹
 Xiaoying Zhou¹
 Edward Smith²
 Francois Haddad¹
 Joseph Wu¹
 Kari Nadeau¹ 

¹Stanford University, Stanford, California

²The Nature Conservancy, San Francisco, California

Correspondence

Mary Prunicki, Stanford University, Stanford, CA.

Email: prunicki@stanford.edu

ORCID

Mary Prunicki  <https://orcid.org/0000-0002-5511-8896>

Kari Nadeau  <https://orcid.org/0000-0002-2146-2955>

REFERENCES

1. D'Amato G, Holgate ST, Pawankar R, et al. Meteorological conditions, climate change, new emerging factors, and asthma and related allergic disorders. A statement of the World Allergy Organization. *World Allergy Organ J.* 2015;8(1):34-37.
2. Reid CE, Brauer M, Johnston FH, Jerrett M, Balmes JR, Elliott CT. Critical review of health impacts of wildfire smoke exposure. *Environ Health Perspect.* 2016;124:1334-1343.
3. Delfino RJ, Brummel S, Wu J, et al. The relationship of respiratory and cardiovascular hospital admissions to the southern California wildfires of 2003. *Occup Environ Med.* 2009;66:189-197.
4. Wettstein ZS, Hoshiko S, Fahimi J, Harrison RJ, Cascio WE, Rappold AG. Cardiovascular and cerebrovascular emergency department visits associated with wildfire smoke exposure in California in 2015. *J Am Heart Assoc.* 2018;11:7.
5. Perera FP, Rauh V, Whyatt RM, et al. Effect of prenatal exposure to airborne polycyclic aromatic hydrocarbons on neurodevelopment in the first 3 years of life among inner-city children. *Environ Health Perspect.* 2006;114:1287-1292.
6. Black C, Gerriets J, Fontaine J, et al. Early life wildfire smoke exposure is associated with immune dysregulation and lung function decrements in adolescence. *Am J Respir Cell Mol Biol.* 2017;56:657-666.
7. Prunicki M, Stell L, Dinakarpandian D, et al. Exposure to NO₂, CO, and PM_{2.5} is linked to regional DNA methylation differences in asthma. *Clin Epigenetics.* 2018;10:2.
8. Nadeau K, McDonald-Hyman C, Noth E, et al. Ambient air pollution impairs regulatory T-cell function in asthma. *J Allergy Clin Immunol.* 2010;126:845-852.
9. Fahy JV. Type 2 inflammation in asthma – present in most, absent in many. *Nat Rev Immunol.* 2015;15:57-65.
10. Olson NC, Sallam R, Doyle MF, Tracy RP, Huber SA. T helper cell polarization in healthy people: implications for cardiovascular disease. *J Cardiovasc Transl Res.* 2013;6:772-786.